

FY2018 Announcement for the influenza vaccination expenses subsidy

【Please make sure to read it before vaccination.】

This vaccination can be taken only if the said person wishes to take it.
There is no obligation of vaccination and it is unnecessary to take it for the person who does not wish to have vaccination as a general rule.

Period From Monday, October 1st, 2018 to Thursday, January 31st, 2019

Eligible person ① Person who has the resident registration in Adachi City and is older than or equal to 65 years old as of the vaccination day.
② Person who has the resident registration in Adachi City, is older than or equal to 60 and under 65 years old and has heart, kidney, or respiratory problem, or immune disorder by HIV comparable to 1st grade of physical disability certificate as of the vaccination day.

※We have sent the vaccination register to the person who is categorized in ① and ②, and who becomes the eligible age too. For the person who becomes eligible on the coming birthday, the vaccination starts from the day before birthday.

※The person who will become eligible age after January 2nd, 2019 is not eligible.

Vaccination fee Copayment is 2,500 yen. Please pay it at the medical institution.

※If the person, who receives the support benefit such as the public assistance recipient and Japanese left behind in China, submit the vaccination register on which “Copayment exempted” is printed, the fee will be **free**.

If the relevant person who possesses the vaccination register on which the amount of copayment is printed, it is necessary to show the certificate of recipient at the Health Prevention Section or each Public Health Center and replace it.

Vaccination place Please refer to the **separate sheet, “List of designated medical institution for FY2018 elderly influenza vaccination”**.

※You can also take vaccination at the designated medical institution in 23 special cities. If you wish to take vaccination in other cities, please make sure the institution is designated in advance.

Vaccination method Please fill in the broad lined box of “**Screening questionnaire and vaccination register of FY2018 elderly influenza vaccination**” and bring it to the designated medical institution.

※Please read carefully the sentence shown below, understand the contents thoroughly and take vaccination when you are in good conditions.

※If the said person who takes vaccination has some paralysis and cannot write the signature on the wish document or cannot confirm the clear resolve because of the dementia, it is absolutely necessary to carefully confirm the resolve for vaccination of the said person by the family member or family doctor and to decide the adaptation of vaccination. If the resolve cannot be confirmed eventually, the vaccination based on the Preventive Vaccination Act cannot be taken.

Before taking vaccination,

(1) Person who cannot take vaccination is

① Clearly to have a high fever.

Generally, it is over 37.5°C of the body temperature.

- ②Clearly to have a serious acute illness.
After the vaccination, because it may become difficult to detect the change in conditions of the person who has an acute illness and is required to take medicine, the vaccination must be postponed as a general rule.
- ③Clearly to have had the anaphylaxis by the component of influenza vaccine before.
“Anaphylaxis” is a severe allergic response occurs within 30 minutes of vaccination. The symptoms of response show the perspiration, acute swelling of face, development of severe rash, nausea, vomiting, hard to vocalize and hard to breathe followed by the shock symptom such as a blood pressure drop.
- ④Clearly to have developed the symptom which suggests the allergic response such as developing high fever, rash and hives occurred within two days after vaccination when the person took the influenza vaccination before.
- ⑤Other cases such that the doctor decides it is an inappropriate.
Even if you do not correspond to ①~④ mentioned above and if the doctor decides the vaccination is inappropriate, you cannot take vaccination.

(2) Person who must consult family doctor when the said person takes vaccination.

- ①The person who clearly has the underlying disease such as a cardiovascular, kidney, liver and blood disease.
- ②The person who has a history of convulsion.
- ③The person who was diagnosed to have immunodeficiency or there is a close relative who has an inborn immunodeficiency disease.
- ④The person who has respiratory disease such as the interstitial pneumonia and bronchial asthma.
- ⑤When you took vaccination before, the symptom which suggests the allergic response such as developing high fever, rash and hives occurred within two days after vaccination.

Precautions for after vaccination

- ①The acute adverse event sometimes occurs within 30 minutes of vaccination. Please make sure that you can contact a doctor or medical institution immediately.
- ②Most of the adverse events of influenza vaccination occur within 24 hours. Please take care of the body conditions carefully during the period.
- ③You can take a bath on the day of vaccination. However, please do not rub the injected part. It is all right to spend the usual life, but please avoid a hard exercise and deep drinking.

Adverse event of vaccination

Although the injection scar is reddened, swollen and pained, developing high fever, chill, headache and whole body lassitude, it will be usually cured in a couple of days. If these symptoms are severe or become prolonged, please take a medical examination by a doctor or medical institution. Beyond that, if you have unclear points, please inquire us.



Inquiry Call Adachi 3880-0039

From 8:00 a.m. to 8:00 p.m. (Every day except form January 1st to 3rd)