FY2026 Notice of COVID-19 Vaccination

【Please read and understand the following carefully and receive the vaccination only when you are in good health.】

This vaccination is available only to individuals who wish to receive it. There is no obligation to receive the vaccination, and individuals who do not wish to receive it are not required to do so.

■**Implementation period**

From October 1 (Thu.), 2026 to March 31 (Wed.), 2027

■**Eligible persons**

① Persons **aged 65 or older** who are registered as residents of Adachi City as of the date of vaccination.

② Persons **aged 60 to under 65** who are registered residents of Adachi City as of the date of vaccination and who have a disability affecting the functions of the heart, kidney, or respiratory organs, or an immune disorder caused by human immunodeficiency virus (HIV). (equivalent to Grade 1 of the Physical Disability Certificate)

※ Persons **who are turning 60 or 65** may receive the vaccination under the Immunization Act starting the day before their birthday. If you wish to receive vaccination before that date, refer to the separate information sheet.

■**Vaccination Fee**

Free of charge (once only during the vaccination period)

※ The vaccine is valued at approximately 16,000 yen.

※ Note that the second and subsequent doses will be subject to a fee (approximately 16,000 yen).

■**Vaccination locations**

Designated medical institutions within Adachi City（refer to the attached list）

※Vaccination is also available at designated medical institutions within Tokyo's 23 special wards.

※Persons aged 59 or 64 may receive vaccination only at the designated medical institutions within Adachi City.

※Details such as the type of vaccine, start date, vaccination days and hours, and whether an appointment is required vary by medical institution. As these details may change during the fiscal year, please be sure to check with the medical institution in advance.

■**How to receive the vaccination**

On the day of vaccination, fill out the section inside the bold-bordered box on the Vaccination Register and Screening Questionnaire and bring it with you to the designated medical institution.

※If you wish to receive the vaccination before reaching the eligible age of 60 or 65, please be sure to bring the attached document titled "For persons who are 59 or 64 years old as of October 2."

※This vaccination is not mandatory. It is administered only if the individual wishes to receive it.

※If you have difficulty signing the form yourself, a family member or your attending physician will make the decision after carefully confirming your wishes. Vaccination cannot be administered if your wishes cannot be confirmed.

What is COVID-19?

(1) How does one become infected?

You can become infected by inhaling viruses in the air released when an infected person coughs, sneezes, or speaks.

(2) What are the main symptoms?

Fever, sore throat, cough, and other symptoms

Tips for preventing infection

To prevent infection, maintain good health by getting adequate nutrition and sleep on a regular basis. When COVID-19 is spreading, avoid crowded places and make sure to wash and disinfect your hands frequently and ventilate your room regularly.

Before receiving vaccination

(1) Things to keep in mind before vaccination

① Make an informed decision before receiving the vaccination.

Please receive the vaccination only after understanding the effects of the vaccine and its adverse reactions. If you have any concerns, consult a doctor beforehand. If you are not satisfied that you understand or agree with the vaccination, you do not need to receive it.

② Fill out the Vaccination Register and Screening Questionnaire accurately.

The vaccination register and screening questionnaire is an important document that allows the doctor to determine whether it is safe for you to receive the vaccination. Please provide accurate information about your health condition.

continue to the back side

(2) Persons who cannot receive the vaccination

① Persons with an obvious fever (generally temperature 37.5℃ or higher)

② Persons with a serious acute illness:
Persons who need to take medication for an acute illness should, as a general rule, postpone vaccination on that day, because it may be difficult to determine how the illness is progressing after vaccination.

③ Persons with a history of severe hypersensitivity, such as anaphylaxis, to any of the ingredients in the vaccine:
Persons who have previously experienced anaphylaxis or multiple symptoms suggestive of anaphylaxis, such as generalized skin or mucosal symptoms, wheezing, difficulty breathing, rapid heartbeat, or a drop in blood pressure

④ Persons for whom a doctor determines that vaccination is otherwise inappropriate

(3) Persons who must consult the attending doctor before receiving the vaccination

① Persons receiving anticoagulant therapy, or those with thrombocytopenia or a coagulation disorder

② Persons who have previously been diagnosed with an immunodeficiency disorder, or who have a close relative with congenital immunodeficiency disorder

③ Persons with underlying conditions such as diseases of the heart, blood vessels, kidneys, liver, or blood, or developmental disorders

④ Persons who, after a previous vaccination, developed symptoms suggestive of an allergic reaction, such as fever or a generalized rash, within two days of vaccination

⑤ Persons with a history of convulsions (seizures)

⑥ Persons who may develop an allergic reaction to any ingredients of this vaccine

(4) Interval between COVID-19 vaccine and other vaccines

The COVID-19 vaccine may be administered at the same time as other vaccines, such as influenza and pneumococcal vaccines, if a doctor specifically determines that concurrent administration is necessary. There are no restrictions on the interval between a COVID-19 vaccine and other vaccinations.

Precautions after vaccination

① Sudden adverse reactions may occur within 30 minutes of vaccination. Please make sure that you can contact a doctor or medical institution promptly if necessary.

② Keep the injection site clean. There is no particular problem with taking a bath on the day of vaccination, but do not rub the injection site firmly.

③ Avoid strenuous exercise and excessive alcohol consumption on the day of vaccination.

Adverse reactions to vaccination

After receiving the vaccine, you may experience pain at the injection site, fatigue, fever, headache, joint pain, or other symptoms. These symptoms usually resolve in a few days. If the symptoms are severe or persist, please consult a doctor or medical institution.
In addition, the person who received the vaccination or their guardian may report symptoms occurring after vaccination to Adachi City using the "Report on Symptoms Occurring After Vaccination" form.

Remedial Measures for Health Damage caused by Vaccination

If you suffer health damage as a result of an adverse reaction to this vaccination, such as a disability that interferes with your daily life, you may be eligible for relief under the Immunization Act in the case of a routine vaccination (including benefits such as medical expenses and disability pensions). In the case of a voluntary vaccination, you may be eligible for relief under the Act on the Pharmaceuticals and Medical Devices Agency.

◆Inquiries◆

Inquiry Call Adachi **3880-0039**

From 8:00 a.m. to 8:00 p.m. (every day except January 1 to 3)

Inquiries: Inoculation Subsection, Health Preventive Control Section, Hygiene Division, TEL 3880-5094

Adachi City